

Parasitic Causes of Pericarditis

JERROLD A. TURNER, MD*

ALTHOUGH MYOCARDITIS is frequent in African trypanosomiasis and American trypanosomiasis (Chagas' disease), involvement of the heart is an uncommon association with other parasitic diseases and pericarditis is even more rare.

Pericardial effusions containing larvae have been described in severe infections with *Trichinella spiralis*, but the myocarditis in these patients overshadows the pericardial involvement. Toxoplasmosis has been reported as a cause of pericarditis.^{67,68} Usually, the diagnosis has been made by serologic methods. One patient with a serologic response diagnostic of toxoplasmosis had pericarditis with progressive pericardial constriction.⁶⁹

Microfilariae have been found in the fluid aspirated from a pericardial effusion,⁷⁰ and constrictive pericarditis has been described in a patient in whom a pericardial effusion associated with filariasis developed.⁷¹

Pericarditis is an uncommon but well described complication of amebic liver abscess. In almost all patients, the abscess has been located in the left lobe of the liver, but there have been rare instances in which abscesses of the right lobe have extended into the pericardial sac.⁷²

Initially, the pericardial involvement may be a serous or serosanguinous effusion (presuppurative stage), but this progresses to a purulent effusion when there is direct communication between the liver abscess and the pericardial cavity (suppurative stage). Although many patients will have a history of travel in areas of the world where amebiasis is common, it should be emphasized that amebic liver abscess and pericarditis occur sporadically in temperate areas, including the United States. The clinical manifestations usually begin with signs and symptoms of a left lobe liver abscess. Epigastric or low retrosternal pain, nausea, vomiting, left shoulder pain, cough, dyspnea, low grade fever, weight loss and tender hepatomegaly are common findings.^{73,74} A few patients present solely with symptoms of pericarditis. Pericardial involvement is frequently

suggested by rapid progression of symptoms and the appearance of signs of pericardial tamponade. Enlargement of the cardiac outline and the "water bottle" configuration of pericardial effusion are commonly seen on chest roentgenograms. Electrocardiography shows changes seen in other forms of pericarditis. Low voltage is also frequently present.

Patients with amebic pericarditis who are misdiagnosed or receive inadequate treatment usually die of pericardial tamponade. When adequate therapy is initiated early, two thirds of the patients may be expected to survive.⁷⁴ A few patients have developed constrictive pericarditis following resolution of the acute problem and in spite of adequate antiamebic therapy.

Diagnosis of amebic pericarditis is dependent on detecting the underlying liver abscess by hepatic scan or echogram. Results of serologic tests for amebiasis have been positive in more than 90 percent of patients with hepatic amebiasis. Examination of material aspirated from the liver or pericardium may reveal trophozoites of *Entamoeba histolytica*. Fecal examination should be carried out but findings are usually negative for amebae.

Immediate treatment is directed toward relieving or preventing pericardial tamponade. This may be done by percutaneous needle aspiration, repeated as frequently as necessary,⁷⁴ or by the creation of a subxiphoid pericardial window under local anesthesia.⁷⁵ Specific antiamebic therapy should be instituted immediately. Metronidazole has become the drug of choice for the treatment of hepatic amebiasis; however, doubt has been raised about its efficacy in patients with pericardial involvement.⁷² Metronidazole attains a high concentration in the liver, but not in the myocardium. Until more information is available, treatment of amebic pericarditis should consist of a combination of chloroquine phosphate 250 mg four times a day for two days then twice a day for 28 days and emetine hydrochloride 1 mg per kg of body weight with 65 mg maximum given intramuscularly daily for 10 days (alternate dihydroemetine dihydrochloride 1 to 1.5 mg per kg of body weight for ten days; maximum total dose 1 gram). A luminal amebicide such as diiodohydroxyquin 650 mg three times a day for 21 days will reduce the likelihood of a persistent, undetectable enteric infection with *E. histolytica*.

Patients should be followed closely after treatment for signs of constrictive pericarditis.

Hydatid disease is caused by the ingestion of

*Assistant Medical Director, Harbor General Hospital.

eggs of the tapeworm *Echinococcus granulosus*. The source of the eggs is usually dog feces, but other carnivores are capable of harboring the adult tapeworm. The ingested eggs hatch in the intestine, and the liberated embryos burrow through the intestinal mucosa and enter the vasculature. They are carried to various tissues where they develop into the larval form, the hydatid cyst. The embryos must pass the filters of the liver and lungs before reaching the blood supply of the myocardium. Therefore, it is not surprising that the majority of patients with hydatid disease have cysts localized to the liver or lungs. Myocardial involvement is seen in only 0.3 to 3.3 percent of cases of hydatid disease.⁷⁶

Pericardial involvement occurs when the cyst in the myocardium ruptures into the pericardial sac. This event frequently is not clinically detectable, or it may lead to a fibrinous or fibrinopurulent pericarditis.⁷⁶ In addition, larval tissue may produce secondary cysts. As the secondary cysts enlarge, they may cause a pericardial reaction and the signs and symptoms of constrictive pericarditis.⁷⁷ An acute pericarditis has been reported in association with allergic manifestations including anaphylaxis.^{78,79} This condition may progress to massive pericardial⁸⁰ effusion with tamponade. A fibrotic reaction of the epicardium which impaired cardiac function and mimicked constrictive pericarditis has also been reported.⁸⁰

The diagnosis of echinococcosis may be suspected in patients who have lived in endemic areas. There may be roentgenographic features of unusual bulges in the cardiac silhouette, intracardiac or pericardial calcifications and, infrequently, the finding of hydatid cysts in the liver or lung. The electrocardiogram may show diminished R waves or the presence of Q waves over the area of the intracardiac cyst. Deep T wave inversion may also be seen in the affected precordial leads.

Skin testing with hydatid fluid (Casoni test) may show reactivity, but false negatives are not uncommon. Serologic tests may be helpful but a negative result does not rule out hydatid disease.

Surgical removal of the cysts is the only effective treatment. Myocardial cysts may be technically very difficult to excise depending upon their location. It is essential that remnants of larval tissue be destroyed or secondary cysts may develop. Formalin, alcohol and 33 percent sodium chloride solution have been used to sterilize cyst cavities. Because of the nature of the myocardium, the concentrated salt solution is considered the

agent of choice.⁷⁷ The use of formalin in the pericardial sac may cause a chemical constrictive pericarditis.

Patients who have had pericardial or myocardial cysts removed require careful follow-up. Electrocardiographic changes may indicate the development of secondary cysts and the need for reoperation.

REFERENCES

1. Spodick DH: Pericardial friction—Characteristics of pericardial rubs in 50 consecutive, prospectively studied patients. *N Engl J Med* 278:1204-1207, May 1968
2. Surawicz B, Lasseter KC: Electrocardiogram in pericarditis. *Am J Cardiol* 26:471-474, Nov 1970
3. Spodick DH: Diagnostic electrocardiographic sequences in acute pericarditis—Significance of PR segment and PR vector changes. *Circulation* 48:575-580, Sep 1973
4. Shabetai R, Fowler NO, Guntheroth WG: The hemodynamics of cardiac tamponade and constrictive pericarditis. *Am J Cardiol* 26:480-489, Nov 1970
5. Craig RJ, Whalen RE, Behar VS, et al: Pressure and volume changes of the left ventricle in acute pericardial tamponade. *Am J Cardiol* 22:65-74, Jul 1968
6. Spodick DH: Acute cardiac tamponade: Pathologic physiology, diagnosis and management. *Progr Cardiovas Dis* 10:64-96, Jul 1967
7. De Cristofaro D, Liu CK: The hemodynamics of cardiac tamponade and blood volume overload in dogs. *Cardiovas Res* 3:292-298, Jul 1969
8. Feigenbaum H: Echocardiographic diagnosis of pericardial effusion. *Am J Cardiol* 26:475-479, Nov 1970
9. Smith WG: Cocksackie B myopericarditis in adults. *Am Heart J* 80:34-36, Jul 1970
10. Cookson H: Virus endocarditis, In Abstracts of papers of Second World Congress of Cardiology. American Heart Association, New York, 1954, p 454
11. DePasquale NP, Burch GE, Shih CS, et al: Experimental Cocksackie virus B₄ valvulitis in cynomolgus monkeys. *Am Heart J* 71:678-683, May 1966
12. Burch GE, Giles TD, Colcolough HL: Pathogenesis of "rheumatic" heart disease—Critique and theory. *Am Heart J* 80:556-561, Oct 1970
13. Robertson R, Arnold CR: Constrictive pericarditis with particular reference to etiology. *Circulation* 26:525-529, Oct 1962
14. Azar GJ: Acute non-specific pericarditis complicated by the development of constrictive pericarditis. *Am Heart J* 65:474-481, Apr 1963
15. Burch GE, Giles TD: The role of viruses in the production of heart disease. *Am J Cardiol* 29:231-240, Feb 1972
16. Lerner AM, Wilson FM: Virus myocardiopathy. *Progr Med Virol* 15:63-91, (Karger, Basel) 1973
17. Sun SC, Sohal RS, Burch GE, et al: Cocksackie virus B₄ pancarditis in cynomolgus monkeys resembling rheumatic heart lesions. *Br J Exp Path* 48:655-661, Dec 1967
18. Gardiner AJS, Short D: Four faces of acute myopericarditis. *Br Heart J* 35:433-442, Apr 1973
19. Sainani GS, Krompotic E, Slodki SJ: Adult heart disease due to the coxsackie virus B infection. *Medicine* 47:133-147, Mar 1968
20. Wilson FM, Miranda QR, Chason JL, et al: Residual pathologic changes following murine coxsackie A and B myocardiitis. *Amer J Path* 55:253-265, May 1969
21. Lennette EH, Shinomoto TT, Schmidt NJ, et al: Observations on the neutralizing antibody response to group B Cocksackie viruses in patients with central nervous system disease. *J Immunol* 86:257-266, Mar 1961
22. Schmidt NJ, Magoffin RL, Lennette EH: Association of group B Cocksackie viruses with cases of pericarditis, myocarditis, or pleurodynia by demonstration of immunoglobulin M antibody. *Inf Imm* 8:341-348, Sep 1973
23. Gatamaitan BG, Chason JL, Lerner AM: Augmentation of the virulence of murine coxsackie virus B-3 myocardiopathy by exercise. *J Exp Med* 131:1121-1136, Jun 1970
24. Kilbourne ED, Wilson CB, Perrier D: The induction of gross myocardial lesions by a Cocksackie (pleurodynia) virus and cortisone. *J Clin Invest* 35:362-370, Apr 1956
25. Voigt GC: Steroid therapy in viral myocarditis. *Am Heart J* 75:575-576, Apr 1968
26. Boyle JD, Pearce ML, Guze LB: Purulent pericarditis—Review of literature and report of 11 cases. *Medicine* 40:119-144, May 1961
27. Varma BN, Smith JM: Tuberculous pericarditis—A review of 17 recent cases. *Tubercle Lond* 48:160-162, 1967

PERICARDITIS

28. Crawford SE, Crook WG, Harrison WW, et al: Histoplasmosis as a cause of acute myocarditis and pericarditis—Report of occurrence in siblings and a review of the literature. *J Pediatr* 28:92-95, Jul 1961
29. Friedman JL, Baum GL, Schwarz J: Primary pulmonary histoplasmosis—Associated pericardial and mediastinal manifestations. *Am J Dis Child* 109:298-303, Apr 1965
30. Gregoriades DG, Langeluttig HV, Polk JW, et al: Pericarditis with massive effusion due to histoplasmosis. *JAMA* 178:331-334, Oct 1961
31. Larson R, Scherb RE: Coccidioidal pericarditis. *Circulation* 7:211-217, 1953
32. Duvall CP, Carbone PP: Cryptococcus neoformans pericarditis associated with Hodgkin's disease. *Ann Intern Med* 64:850-856, Apr 1966
33. Herman GR, Marchand EJ, Greer GH, et al: Pericarditis—Clinical and laboratory data of 130 cases. *Am Heart J* 43:641-652, May 1952
34. Pereira MD: Actinomycosis of the pericardium. *Med Ann DC* 29:28-33, Jan 1960
35. Chavez CM, Causey WA, Conn JH: Constrictive pericarditis due to infection with *Nocardia asteroides*. *Chest* 61:79-81, Jan 1972
36. Susens GP, Al-Shamma A, Rowe JC, et al: Purulent constrictive pericarditis caused by *Nocardia asteroides*. *Ann Intern Med* 67:1021-1032, Nov 1967
37. Fraumeni JF, Fear RE: Purulent pericarditis in aspergillosis. *Ann Intern Med* 57:823-828, Nov 1962
38. Connolly DC, Burchell HB: Pericarditis—A ten year survey. *Am J Cardiol* 7:7-14, Jan 1961
39. Rooney JJ, Croces JA, Lyons HA: Tuberculosis pericarditis. *Ann Intern Med* 72:73-78, Jan 1970
40. Robertson R, Arnold CR: Constrictive pericarditis with particular reference to etiology. *Circulation* 26:525-529, Oct 1962
41. Mynors JM, Reichman B: Pericarditis, a five year study in the African. *Cent Afr J Med* 19:19-22, Feb 1973
42. Kaplan MM, Sherwood LM: Acute pericarditis due to histoplasma capsulatum. *Ann Intern Med* 58:862-867, May 1963
43. Klieger HL, Fisher ER: Fibrocalcific constrictive pericarditis due to histoplasma capsulatum. *N Engl J Med* 267:593-596, Sep 1962
44. Saslaw S, Norfleet RG: Acute histoplasma pericarditis. *Arch Intern Med* 122:162-166, Aug 1968
45. DeRoover P, Lacquet A: Pericarditis following pneumonectomy for pulmonary tuberculosis. *Br J Dis Chest* 53:356-359, Oct 1959
46. Shugoll GI: Percutaneous myocardial and pericardial biopsy with the Menghini needle. *Am Heart J* 85:35-39, Jan 1973
47. Scientific Assembly statement—The use of skin tests and serologic tests in histoplasmosis, coccidioidomycosis, and blastomycosis. *Am Rev Resp Dis* 108:156-159, Jul 1973
48. Heiner DC: Diagnosis of histoplasmosis using precipitin reactions in agar gel. *Pediatrics* 22:616-627, Oct 1958
49. Wychulis AR, Connolly DC, McGoon DC, et al: Surgical treatment of pericarditis. *J Thorac Cardiovasc Surg* 62:608-617, Oct 1971
50. Hageman JH, D'Esopo ND, Glenn WW: Tuberculosis of the pericardium—A long-term analysis of 44 proved cases. *N Engl J Med* 270:327-332, Feb 1964
51. Schepers GWH: Tuberculous pericarditis—Review. *Am J Cardiol* 9:248-276, Feb 1962
52. Portal RW, Besterman EMM, Chambers RJ, et al: Prognosis after operation for constrictive pericarditis. *Br Med J* 1:563-569, Mar 1966
53. Simcha A, Taylor JFN: Constrictive pericarditis in childhood. *Arch Dis Child* 46:515-519, Aug 1971
54. Somers K, De Buse PJ, Patel AK, et al: Childhood tuberculous pericarditis. *Chest* 60:22-28, Jul 1971
55. Case records of the Massachusetts General Hospital, constrictive tuberculous pericarditis, case 30—1972. *N Engl J Med* 287:189-193, Jul 1972
56. Logue B, Tutunji F: The use of steroids in pericarditis. *Am Heart J* 64:570-571, Oct 1962
57. Flory WD: Pericarditis—Report of seven cases. *Minn Med* 45:502-510, May 1962
58. Still GF: Observations on suppurative pericarditis in children. *Br Med J* 2:606-608, Sep 1901
59. Kauffman CA, Watanakunakorn C, Phair JP: Purulent pneumococcal pericarditis—A continuing problem in the antibiotic era. *Am J Med* 54:743-750, Jun 1973
60. Kikolaew GF: Du role des abcès du myocarde dans la pathogène des pericardites suppurees. *Presse Med* 45:926-928, Jun 19, 1937
61. Rose RL, Higgins LS, Helgason AH: Bacterial endocarditis, pericarditis and cardiac tamponade. *Am J Cardiol* 19:447-451, Mar 1967
62. Buchbinder NA, Roberts WC: Left-sided valvular active infective endocarditis—A study of forty-five necropsy patients. *Am J Med* 53:20-36, Jul 1972
63. Beck D: Pericarditis and subacute bacterial endocarditis. *J Mt Sinai Hosp* 8:364-375, 1942
64. Gersony WM, McCracken GH Jr: Purulent pericarditis in infancy. *Pediatrics* 40:224-232, Aug 1967
65. Levin HS, Hosier DM: Salmonella pericarditis—Report of a case and review of the literature. *Ann Intern Med* 55:817-823, Nov 1961
66. Scott LP, Knox D, Perry LW, et al: Meningococcal pericarditis—Report of 2 Cases—I. Complicated by acute constrictive pericarditis. *Am J Cardiol* 29:104-108, Jan 1972
67. Hakilla J, Frick HM, Halonen PI: Pericarditis and myocarditis caused by toxoplasma—Report of a case and review of the literature. *Am Heart J* 55:758-765, May 1958
68. Jones TC, Kean BH, Kimball AC: Pericarditis associated with toxoplasmosis—Report of a case and review of the literature. *Ann Intern Med* 62:786-790, Apr 1965
69. Nagaratnam N, Stephen SJ: Chronic constrictive pericarditis associated with toxoplasmosis. *Br Heart J* 35:868-870, Aug 1973
70. Foster DG: Filariasis—A rare cause of pericarditis. *J Trop Med Hyg* 59:212-214, 1956
71. Charan A, Sinha K: Constrictive pericarditis following filarial effusion. *Indian Heart J* 25:213-215, Jul 1973
72. Watson RB, Steel RK, Spiegel TM: Amebic pericarditis consequent to amebic abscess of right lobe of the liver. *Am J Trop Med Hyg* 21:889-893, Nov 1972
73. Alkan WJ, Kalmi B, Kalderon M: The clinical syndrome of amebic abscess of the left lobe of the liver. *Ann Intern Med* 55:800-813, Nov 1961
74. MacLeod IN, Wilmot AJ, Powell SJ: Amoebic pericarditis. *Quart J Med* 35:293-311, Apr 1966
75. Fontenelle LJ, Cuella L, Dooley BN: Subxiphoid pericardial window. *J Thor Cardiovasc Surg* 62:95-97, Jul 1971
76. Kean BH, Breslau RC: Parasites of the Human Heart. New York, Grune and Stratton, 1964
77. Pinch LW, Sehat A: Constrictive pericarditis associated with echinococcus cyst. *Chest* 61:194-195, Feb 1972
78. Perez-Gomez F, Duran H, Tamames S, et al: Cardiac echinococcosis—Clinical picture and complications. *Br Heart J* 35:1326-1331, Dec 1973
79. DiBello R, Mantero ME, Dubra J, et al: Hydatid cyst of the left ventricle of the heart. *Am J Cardiol* 19:603-606, Apr 1967
80. DiBello R, Abo JC, Borges UL: Hydatid constrictive pericarditis. *J Thorac Cardiovasc Surg* 59:530-532, Apr 1970